PROPAGATION

PELARGONIUM SARITA®

Pelargonium x hybrida

TRAY & ROOTING

Form **Vegetative Cutting**

Rooting Hormone **Yes**

Tray Size **32, 52, 72, 102 strip**

Plants Per Cell 1

Days In Mist 8-16

Wks. From Stick To

Transplant 3-4

Pinch? No

Days To Pinch N/A

FERTILITY

Fertilize Through Mist? No

Fertilizer Rate Moderate: 100-150 ppm

(20-10-20 constant feed)

EC Range Moderate (1.0-1.5 mS via SME)

pH Range **6.0-6.4**

TEMPERATURE

Vernalization Required? **No**

Average Soil

Temperature

68-72°F (20-22°C)

Average Air

Temperature **65-72°F (18-22°C)**

IRRIGATION FREQUENCY

Level 3: keep media moderately moist at all times

LIGHT (SUPPLEMENTAL)

No

SHADE

Keep light intensity between 700-1200 fc (7500-12900 lux) until callus formation, 2000-2500 fc (21500-26900 lux) during root initiation, and 2500-4000 fc (26900-43100 lux) for toning.

POTENTIAL DISEASES

Xanthomonas, Rhizoctonia, Botrytis

POTENTIAL PESTS

thrips, aphids, fungus gnats

PGR SUGGESTIONS

- 1,500-2500 ppm *Daminozide* (85%) spray
- Ethephon (3.9%) at 350-500 ppm can be used for bud removal and branching 12-21 days after stick (when liners are rooted). Do not apply within 6-8 weeks of ship date.





Sarita® Fire

TIPS

 Apply a 200 ppm IBA spray (or equivalent product) within 24 hours of sticking to aid root formation.

WE'RE HERE TO HELP YOU GROW

Tech.NA@DummenOrange.com 614-850-9551

DummenOrange.com

FINISHING

PELARGONIUM SARITA®

Pelargonium x hybrida

CONTAINER SIZE	PLANTS PER POT		MID SEASON	LATE SEASON
6 in. (15 cm)	1	6-8	5-7	4-6
8 in. (19 cm)	1-2	7-11	7-10	7-9
10 in. (22 cm)	2-3	8-12	8-11	8-10
12 in. (30 cm)	3-4	8-12	8-11	8-10

	75	31	1	τ	
Εŀ	~ 1				V
	\ L				ш

Moderate: 125-200 ppm N
(15-5-15 at or similar calciumnitrate formula alternated with
20-10-20)

EC Range High (2.0-2.5 mS via SME)

pH Range **6.0-6.4**

TEMPERATURE

Day Temperature **65-75°F (18-24°C)**

Night Temperature **58-62° (14-17°C)**

IRRIGATION

Irrigation Frequency **Level 3: keep media moderately moist at all times**

LIGHT

Daylength Neutral? Yes

Supplemental No

Benefits From Shade? No

PINCHING

Finishing Pinch? **No**

POTENTIAL DISEASES

Xanthomonas, Rhizoctonia, Botrytis

POTENTIAL PESTS

thrips, aphids, fungus gnats

PGR SUGGESTIONS

- 750-1500 ppm *Chlormequat* (11.8%) as needed
- 1200 ppm *Chlormequat* (11.8%) + 2500 ppm *Daminozide* (85%) tank mix





Sarita® Neon Sizzle

TIPS

- Cultivars in this series are relatively vigorous and will likely require PGR applications.
- Utilize fertilizer formulations low in phosphorous and ammonium to help control stretch, increase branching, increase shelf life, and encourage full flowering.
- Consider direct-stick with callused cuttings to reduce overall production time by 1-2 weeks.

WE'RE HERE TO HELP YOU GROW

Tech.NA@DummenOrange.com 614-850-9551

DummenOrange.com

FEATURES AND BENEFITS

PELARGONIUM SARITA®

Pelargonium x hybrida

DÜMMEN ORANGE

DESCRIPTION

Sarita® is a medium vigor dark foliaged interspecific series with large single flowers.

KEY GARDEN INFORMATION

Exposure Sun to Partial Shade

Height **12-18 inches (30-46cm)**

Garden Spacing 12-18 inches (30-46cm)

Watering Allow soil to dry slightly instructions between thorough waterings.

Hardiness Zone 10-12

CONSUMER APPEAL

• Interspecifics are known for their heat tolerance and versatility, this series is no exception.

GARDEN ATTRIBUTES

- If planted in the landscape, *Pelargonium* prefer rich well-drained soils. However, they can tolerate heavy clay soils if not overwatered.
- Although not favored by bees, the long bloom season and bright flowers of *Pelargonium* naturally attract hummingbirds. Additionally, their recognizable foliage aroma is a natural deterrent to rabbits, dear, and herbivorous insects.
- *Pelargonium* can be overwintered indoors for extended seasonal interest.



Sarita® Wild Salmon

NATIVITY

Of hybrid origin: parental lineages native to Southern Africa.

WE'RE HERE TO HELP YOU GROW

Tech.NA@DummenOrange.com 614-850-9551 DummenOrange.com